

REPRESENTATION THEORY FOR THE KRIŽ MODEL

SAMIA ASHRAF¹, HANIYA AZAM², BARBU BERCEANU³

ABSTRACT. The natural action of the symmetric group on the configuration spaces $F(X, n)$ induces an action on the Križ model $E(X, n)$. The representation theory of this DGA is studied and a big acyclic subcomplex which is S_n -invariant is described.

1. INTRODUCTION

The ordered configuration space of n points $F(X, n)$ of a topological space X is defined as

$$F(X, n) = \{(x_1, x_2, \dots, x_n) \in X^n \mid x_i \neq x_j \text{ for } i \neq j\}.$$

For X a smooth complex projective variety, I. Križ [K] constructed a rational model $E(X, n)$ for $F(X, n)$, a simplified version of the Fulton-MacPherson model [FM]. Recently Lambrechts and Stanley [LaSt] constructed a (quasi)-model for the configuration space of a topological space with Poincaré duality cohomology; if such a space is formal, the model of Lambrechts-Stanley is reduced to the Križ model and this is the case of Kähler manifolds, see [DGMS]. Therefore all the results of this paper could be applied to (simply connected) formal closed manifolds (with few changes for the odd-dimensional manifolds).

Let us remind the construction of Križ. We denote by $p_i^* : H^*(X) \rightarrow H^*(X^n)$ and $p_{ij}^* : H^*(X^2) \rightarrow H^*(X^n)$ (for $i \neq j$) the pullbacks of the obvious projections and by m the complex dimension of X (for cohomology groups we use rational or complex coefficients). The model $E(X, n)$ is defined as follows: as an algebra $E(X, n)$ is isomorphic to the exterior algebra with generators G_{ij} , $1 \leq i, j \leq n$ (of degree $2m - 1$) and coefficients in $H^*(X)^{\otimes n}$ modulo the relations

$$\begin{aligned} G_{ji} &= G_{ij} \\ p_j^*(x)G_{ij} &= p_i^*(x)G_{ij}, & (i < j), x \in H^*(X) \\ G_{ik}G_{jk} &= G_{ij}G_{jk} - G_{ij}G_{ik}, & (i < j < k). \end{aligned}$$

The differential d is given by $d|_{H^*(X)^{\otimes n}} = 0$ and $d(G_{ij}) = p_{ij}^*(\Delta)$, where $\Delta = w \otimes 1 + \dots + 1 \otimes w \in H^*(X) \otimes H^*(X)$ denotes the class of the diagonal and $w \in H^{2m}(X)$ is the fundamental class.

This model is a differential bigraded algebra $E(X, n) = \bigoplus_{k,q} E_q^k(X, n)$: the lower degree q (called the exterior degree) is given by the number of exterior generators

Key words and phrases. Representations of symmetric groups, configuration spaces, rational model.

2010 AMS Classification Primary: 55R80, 20C30; Secondary: 55P62, 13A50.

This research is partially supported by Higher Education Commission, Pakistan.

G_{ij} , and the upper degree k is given by the total degree: the multiplication is homogeneous

$$E_q^k \otimes E_{q'}^{k'} \rightarrow E_{q+q'}^{k+k'}$$

and the differential has bidegree $\begin{pmatrix} +1 \\ -1 \end{pmatrix}$

$$d : E_q^k \rightarrow E_{q-1}^{k+1}.$$

In the next definition $G_{I_* J_*}$ is a product of exterior generators


$$G_{I_* J_*} = G_{i_1 j_1} G_{i_2 j_2} \cdots G_{i_q j_q}.$$

Definition 1.1. (Fulton-MacPherson [FM], Križ [K]) *The symmetric group $\mathcal{S}_n$ acts on $E(X, n)$ by permuting the factors in $H^*(X^n) = H^{*\otimes n}$ and changing the indices of the exterior generators: for an arbitrary permutation $\sigma \in \mathcal{S}_n$*

$$\sigma(p_1^*(x_{h_1}) \cdots p_n^*(x_{h_n}) G_{I_* J_*}) = p_{\sigma(1)}^*(x_{h_1}) \cdots p_{\sigma(n)}^*(x_{h_n}) G_{\sigma(I_*) \sigma(J_*)}.$$

The action of $\mathcal{S}_n$ is well defined because the set of relations is invariant under this action.

In the next diagram the nonzero bigraded components $E_q^k(X, n)$ lie in the trapezoid with vertices $(0, 0)$, $(2mn, 0)$, $((n-1)(2m-1), n-1)$ and $(n(2m-1)+1, n-1)$.



Along the horizontal lines we describe a duality preserving the symmetric action: the dotted median line of the trapezoid is the axis of symmetry for this duality, see Proposition 2.2 in Section 2. In the same section we introduce the combinatorial “types” of monomials of $E_*(X, n)$: these are parameterized by forests in which every tree contains a cohomology class of X . The type decomposition of the bigraded components gives a direct sum of $\mathcal{S}_n$ submodules, each of these being generated by a unique element: see Theorem 2.13. In the next section we describe the $\mathcal{S}_n$ structure of types: explicit decomposition into irreducible representations in many particular cases and, using the results of Lehrer-Solomon [LeSo], we compute the character for the general type. See the Propositions 3.3, 3.4, 3.8, 3.12 and Theorem 3.15.

In Section 4 we present some properties of the differential which are consequences of its $\mathcal{S}_n$ -equivariance. For all complex projective manifolds, except the projective line, we show that the differential is injective on the “left side” of the trapezoid:

Proposition 1.2. *The differentials in the Križ model of a projective manifold different from $\mathbb{CP}^1$ are injective for any q in the interval $[1, n-1]$:*

$$d : E_q^{q(2m-1)}(X, n) \hookrightarrow E_{q-1}^{q(2m-1)+1}(X, n).$$

The top horizontal line has no contribution to the cohomology, too:

Proposition 1.3. *The top differentials in the Križ model are injective for any k in the interval $[(n-1)(2m-1), n(2m-1)+1]$:*

$$d : E_{n-1}^k(X, n) \hookrightarrow E_{n-2}^{k+1}(X, n).$$

In Section 5 we show that the “right side” of the trapezoid is an acyclic complex:

Proposition 1.4. *All cohomology groups of the subcomplex*

$$0 \rightarrow E_{n-1}^{n(2m-1)+1}(X, n) \rightarrow E_{n-2}^{n(2m-1)+2}(X, n) \rightarrow \dots \rightarrow E_0^{2nm}(X, n) \rightarrow 0$$

are zero.

Other (smaller) copies of this subcomplex are contained in the interior of the trapezoid and their sum gives a large acyclic complex which is also $\mathcal{S}_n$ -equivariant. This subcomplex $E_*^*(w(X, n))$ and the quotient $E_*^*(X, n)/E_*^*(w(X, n))$ are described in Proposition 5.7 and Proposition 5.8; the location of the subcomplex $E_*^*(w(X, n))$ is given in the diagram by the interior lines with slope -1 . In [BMP] a different acyclic subcomplex of $E_*^*(X, n)$ is described: this is an ideal, but is not an $\mathcal{S}_n$ -submodule. The subcomplex $E_*^*(w(X, n))$ is an $\mathcal{S}_n$ -subalgebra, but not an ideal. The right side of the trapezoid, denoted by $E_*^{Top}(X, n)$ in Section 5, is an acyclic ideal which is also an $\mathcal{S}_n$ -submodule, but it is quite small.

In the last section the simplest and, from the viewpoint of Proposition 1.2, the exceptional case of $\mathbb{CP}^1$ is analyzed; we recover and we complete the results of Cohen-Taylor [CT1], [CT2] and Feichtner-Ziegler [FZ], computing in this case the Poincaré polynomials in two variables


$$P_{F(X, n)}(t, s) = \sum_{k, q \geq 0} (\dim H_q^k) t^k s^q.$$

Theorem 1.5. *In the cohomology algebra of the configuration space $F(\mathbb{CP}^1, n)$ ($n \geq 4$) the non zero bigraded components are*

$$H_q^k \cong H_{q+1}^{k+3} \text{ for } q = 0, 1, \dots, n-3.$$

Its Poincaré polynomial is

$$P_{F(\mathbb{CP}^1, n)}(t, s) = (1 + st^3)(1 + 2st)(1 + 3st) \dots (1 + (n-2)st).$$



Other extensions and applications of the results of this paper could be found in [AsB] and [AzB].

For the irreducible $\mathcal{S}_n$ -modules we will use the standard notation (see [FH]): $V(\lambda)$ corresponds to the partition of n , $\lambda \vdash n$, $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_t \geq 1)$, and also the stable notation (see [CF] or [AAB]): $V(\mu)_n = V(n - \sum \mu_i, \mu_1, \mu_2, \dots, \mu_s)$ for $\mu = (\mu_1 \geq \mu_2 \geq \dots \geq \mu_s \geq 1)$ satisfying the relation $n - \sum_{i=1}^s \mu_i \geq \mu_1$.

2. COHOMOLOGY CLASSES IN THE FOREST

We fix a (monomial) ordered basis for the cohomology algebra $H^*(X; \mathbb{Q}) : \mathcal{B} = \{x_1 = 1 \prec x_2 \prec \dots \prec x_B = w\}$, where $w \in H^{2m}(X; \mathbb{Q})$ is the fundamental class of X and $B = \sum \beta_i$ is the sum of Betti numbers; we choose the order $\prec$ such that the sequence $\{\deg x_i\}_{i=1, B}$ is increasing (not necessarily strictly increasing). Using simple computations with Diamond lemma (see [Bg]) one can find a monomial basis for the Križ model: we denote by $G_{I_* J_*} = G_{i_1 j_1} G_{i_2 j_2} \dots G_{i_q j_q}$ the exterior monomial corresponding to the sequences $I_* = (i_1, \dots, i_q)$, $J_* = (j_1, \dots, j_q)$, where $i_a < j_a$ ($a = 1, 2, \dots, q$) and $j_1 < j_2 < \dots < j_q$, and by $x_{H_*} = x_{h_1} \otimes x_{h_2} \otimes \dots \otimes x_{h_n}$ ($x_{h_a} \in \mathcal{B}$) a scalar from $H^{*\otimes n}$. Then

$$\{x_{H_*} G_{I_* J_*} \mid x_{h_a} = 1 \text{ if } a \in J_*, \deg x_{H_*} = k - q(2m - 1)\}$$

is a basis of $E_q^k(X, n)$ and we call it the *canonical* (Bezrukavnikov) basis (see [Bz]).

The next result is obvious:

Proposition 2.1. *The bigraded components $E_q^k(X, n)$ are invariant under the action of the symmetric group and the differential d is $\mathcal{S}_n$ -equivariant:*

$$d(\sigma(x_{H_*} G_{I_* J_*})) = \sigma(d(x_{H_*} G_{I_* J_*})).$$

This proposition and the Schur lemma give a splitting of the Križ complex into subcomplexes corresponding to the decomposition of E_*^* into isotypical components $E_*^*(V(\lambda))$, for λ an arbitrary partition of n :

$$(E_*^*(X, n), d) = \bigoplus_{\lambda \vdash n} (E_*^*(V(\lambda)), d_\lambda).$$

The cohomology algebra $H^*(X; \mathbb{Q})$ satisfies Poincaré duality; denote by $\mathcal{B}^*$ the Poincaré dual basis

$$\mathcal{B}^* = \{y^1 = w, y^2, \dots, y^B = 1 \mid x_i y^j = \delta_{ij} w\}.$$

Proposition 2.2. *For any $q = 0, 1, \dots, n - 1$ and any k in the interval of integers $[(2m - 1)q, 2mn - q]$, there is an isomorphism of $\mathcal{S}_n$ -modules*

$$E_q^k(X, n) \cong E_q^{2mn + 2q(m-1) - k}(X, n).$$

Proof. Define the map $\Phi : E_q^k \rightarrow E_q^{2mn + 2q(m-1) - k}$ on the basis by

$$\Phi(x_{H_*} G_{I_* J_*}) = x'_{H_*} G_{I_* J_*},$$

where the factors of $x'_{H_*} = x'_{h_1} \otimes x'_{h_2} \otimes \dots \otimes x'_{h_n}$ are given by

$$x'_{h_a} = \begin{cases} 1 (= x_{h_a}) & \text{if } h_a \in J_* \\ y^{h_a} & \text{if } h_a \text{ is not in } J_*. \end{cases}$$

It is easy to see that Φ is $\mathcal{S}_n$ -equivariant and the sum of the total degree of $x_{H_*} G_{I_* J_*}$ and the total degree of $\Phi(x_{H_*} G_{I_* J_*})$ is $2mn + 2q(m - 1)$. $\square$


Now we associate to monomials in the Križ model $E_*^*(X, n)$ *marked graphs*; these are extensions of the graphs introduced in [LeSo] in order to study the representation theory of the Arnold algebra, the cohomology algebra of $F(\mathbb{C}, n)$, see [A]. To any monomial $G_{I_* J_*}$ from the Arnold algebra $\mathcal{A}(n)$, Lehrer and Solomon associated a graph Γ with vertices $\{1, 2, \dots, n\}$ and edges $\{i, j\}$ corresponding to the factors G_{ij} of the given monomial. We associate to the monomial $\mu = x_{h_1} \dots \otimes x_{h_n} G_{I_* J_*}$ from $H^*(X)^{\otimes n} \otimes \mathcal{A}(n)$ the graph Γ corresponding to $G_{I_* J_*}$, marking the vertices

$1, 2, \dots, n$ with the monomials $x_{h_1}, \dots, x_{h_n}$ from the fixed basis $\mathcal{B}$. Due to the relation $p_j^*(x)G_{ij} = p_i^*(x)G_{ij}$, in the image of μ in the Križ algebra $E_*(X, n)$ we can move all the factors x_{h_i} corresponding to a given connected component of Γ on the smallest index of that component; in this way the marks of the associated graph of the monomial $\mu = x_{h_1} \dots \otimes x_{h_n} G_{I_* J_*}$ in $E_*(X, n)$ ($x_h = 1$ if $h \in J_*$) are the marks of the connected components of the graph Γ .

Example 2.3. Consider the monomial $\mu = x_{H_*} G_{I_* J_*} \in E_6^*(X, 11)$ given by

$$x_{h_1} \otimes 1 \otimes 1 \otimes x_{h_4} \otimes 1 \otimes 1 \otimes 1 \otimes x_{h_8} \otimes 1 \otimes x_{h_{10}} \otimes x_{h_{11}} G_{12} G_{23} G_{45} G_{46} G_{47} G_{89}.$$

Its associated marked graph is



It is obvious how to obtain an element in $E_*(X, n)$ starting with a graph with vertices $\{1, 2, \dots, n\}$ and marks from $H^*(X)$ on the connected components of this graph.

Remark 2.4. [LeSo] If a marked graph contains a cycle, the associated element in $E_*(X, n)$ is zero. Therefore we will consider only marked forests (all the connected components are trees).

Proof. Start an induction on the length of the cycle with length 3:

$$G_{ij} G_{ik} G_{jk} = G_{ij} (G_{ij} G_{jk} - G_{ij} G_{ik}) = 0.$$

To a cycle of length $l + 1$ corresponds an element containing as a factor the product

$$G_{i_1 i_2} G_{i_2 i_3} \dots G_{i_l i_{l+1}} G_{i_{l+1} i_1} = G_{i_1 i_2} \dots G_{i_{l-1} i_l} (G_{i_1 i_l} G_{i_l i_{l+1}} - G_{i_1 i_l} G_{i_l i_{l+1}}),$$

both terms having associated graphs with cycles of length l . □

If we restrict the correspondence $\{\text{monomials in } E_*(X, n)\} \rightarrow \{\text{marked graphs}\}$ to the canonical (Bezrukavnikov) basis, we obtain only marked monotonic graphs:

Definition 2.5. A tree with vertices $\{1 \leq i_1 < i_2 < \dots < i_p \leq n\}$ is monotonic if, for any vertex i_k , the unique path from i_1 to i_k is strictly increasing:

$$i_1 < i_a < i_b < \dots < i_k$$

(choosing the root i_1 , the rooted tree is monotonic). A forest with vertices $\{1, 2, \dots, n\}$ is monotonic if all its trees are monotonic.

Example 2.6. The tree

$$\underline{3 \quad 2 \quad 5} \text{ is monotonic but } \underline{3 \quad 5 \quad 2} \text{ is not.}$$

Remark 2.7. There is one to one correspondence

$$\{\text{monomials in the canonical basis of } E_*(X, n)\} \leftrightarrow \{\text{marked monotonic forests}\}.$$


Proof. Let us suppose that the graph Γ associated to a canonical monomial $G_{I_* J_*}$ ($j_1 < j_2 < \dots < j_q, i_a < j_a$ for $a = 1, 2, \dots, q$) is connected; from its Euler characteristic we find that $\text{card}(I_* \cup J_*) = q + 1$. If Γ is not monotonic, there is a path $i_1 - \dots - j - k - h$ such that $(i_1 \leq) j < k > h$, and this corresponds to a forbidden product $G_{jk} G_{hk}$ in $G_{I_* J_*}$. Conversely, to any monotonic tree (or forest) corresponds a product $G_{I_* J_*}$ from the canonical basis: a vertex j , distinct from the minimal vertex i in the same connected component, is joined with a unique vertex h , and this is smaller than j , namely the second last vertex on the path from i to j ; therefore j appears only once on the second position, hence in the sequence J_* . $\square$

There is an obvious action of the symmetric group $\mathcal{S}_n$ on the set of marked graphs: the natural action of $\mathcal{S}_n$ on the set of vertices $\{1, 2, \dots, n\}$ induces an action on the set of edges and an action on the connected components and the corresponding marks. The set of monotonic marked forests is not $\mathcal{S}_n$ -stable, like the set of monomials in the canonical basis of the Kriz model; but $E_*(X, n)$ and the $\mathbb{Q}$ vector space generated by marked monotonic forests are $\mathcal{S}_n$ -stable and these two vector spaces will be identified.


Example 2.8. In this example and the next one two $\mathcal{S}_4$ -orbits in the $\mathbb{Q}$ -span of monotonic marked forests are described: $\oplus$ and $\ominus$ stand for the sum and difference in this vector space and $\tau_i = (i, i+1)$ ($i = 1, 2, 3$) are the Coxeter generators of $\mathcal{S}_4$. To save space, the bullet $\bullet$ corresponds to the vertex 1, the root of the tree, and the vertices connected to 1 are, from left to right, written in increasing order; hence



But the next rooted tree is ambiguous therefore we will write




For the same reason the (unique) mark $x_h \in \mathcal{B}$ is omitted.



In the first example we have a “small” orbit: the dimension of the representation is 3 and it is isomorphic to $V(2, 1, 1)$. In the second one we will see a “complete” orbit: the dimension is 6 and the representation is $V(3, 1) \oplus V(2, 1, 1)$.

Example 2.9.



The second example suggests the next definition

Definition 2.10. We say that two monotonic marked forests are of the same type, $(\Gamma, H_*) \sim (\Gamma', H'_*)$, if there is a permutation $\sigma \in S_n$ which induces a bijection between the connected components of the two graphs, preserving the number of elements of the corresponding components and their marks.

It is clear that marked monotonic forests in the same S_n -orbit are of the same type, but not conversely. A complete system of representatives for this equivalence relation is given by forests of bamboos:

$$(\Gamma_{L_*}, H_*) : \quad \begin{array}{ccccccc} & x_{h_1} & & x_{h_2} & & \dots & x_{h_t} \\ \hline 1 & 2 & \dots & L_1 & L_1 + 1 & \dots & L_2 & \dots & L_{t-1} + 1 & \dots & L_t \end{array}$$

where the sequence of lengths of the bamboos $L_* = (\lambda_1, \lambda_2, \dots, \lambda_t)$ is decreasing $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_t$, $L_i = \lambda_1 + \lambda_2 + \dots + \lambda_i$, $L_t = n$, and, for equal lengths, the marks are in a decreasing order. We split the $\mathcal{S}_n$ -modules $E_q^k(X, n)$ into smaller pieces using the type of the associated monotonic marked forests.

Definition 2.11. For a given marked forest of bamboos (Γ_{L_*}, H_*) we define the subspace of type (L_*, H_*) as the linear span of monomials with the associated marked graph of type (Γ_{L_*}, H_*) ; this will be denoted by $E_*^k(L_*, H_*)$.

Example 2.12. For the type $(\Gamma_{(3,1,1)}, (h_1, h_2, h_3))$, where $x_{h_2} \succ x_{h_3}$,

$$(\Gamma_{L_*}, H_*) : \quad \begin{array}{ccccc} & x_{h_1} & & x_{h_2} & x_{h_3} \\ & \bullet & & \bullet & \bullet \\ 1 & \text{---} & 2 & 3 & 4 & 5 \end{array}$$

the associated space $E_*^k(L_*, H_*)$ is of dimension 40 and its canonical basis is given by the monomials

$$\begin{aligned} & x_{h_1} \otimes 1 \otimes 1 \otimes x_{h_2} \otimes x_{h_3} G_{12} G_{13}, & x_{h_1} \otimes 1 \otimes 1 \otimes x_{h_3} \otimes x_{h_2} G_{12} G_{13}, \\ & x_{h_1} \otimes 1 \otimes 1 \otimes x_{h_2} \otimes x_{h_3} G_{12} G_{23}, & x_{h_1} \otimes 1 \otimes 1 \otimes x_{h_3} \otimes x_{h_2} G_{12} G_{23}, \\ & \dots & \dots \\ & x_{h_2} \otimes x_{h_3} \otimes x_{h_1} \otimes 1 \otimes 1 G_{34} G_{45}, & x_{h_3} \otimes x_{h_2} \otimes x_{h_1} \otimes 1 \otimes 1 G_{34} G_{45}. \end{aligned}$$

If in this example $x_{h_2} = x_{h_3}$, the dimension of $E_*^k(L_*, H_*)$ is 20.

To a given type (L_*, H_*) , $L_* = (\lambda_1, \dots, \lambda_t)$, $H_* = (h_1, \dots, h_t)$, we will associate two integers:

$$|L_*| = \sum_{i=1}^t (\lambda_i - 1) \quad \text{and} \quad |H_*| = \sum_{h=1}^t \deg(x_h).$$

Theorem 2.13. The bihomogenous components $E_q^k(X, n)$ can be decomposed into a direct sum of monogenic $\mathcal{S}_n$ -submodules

$$E_q^k(X, n) = \bigoplus_{\substack{|L_*|=q \\ |H_*|=k-q(2m-1)}} E_q^k(L_*, H_*)$$

In particular, the multiplicities of the irreducible $\mathcal{S}_n$ -submodules of each term $E_q^k(L_*, H_*) = \bigoplus_{\lambda \vdash n} m_\lambda V(\lambda)$ satisfy the relations $m_\lambda \leq \dim V(\lambda)$.

Proof. The $\mathcal{S}_n$ -module $E_q^k(X, n)$ is the direct sum $\bigoplus E_q^k(L_*, H_*)$ by the very definition of monomials of type (L_*, H_*) . We will show that:

- a) for any type (L_*, H_*) , the vector space $E(L_*, H_*)$ is $\mathcal{S}_n$ -stable;
- b) a generator of the $\mathcal{S}_n$ -module $E_*^k(L_*, H_*)$ is the monomial corresponding to the marked monotonic bamboo (L_*, H_*) :

$$\mu_{(L_*, H_*)} = p_1^*(x_{h_1}) p_{L_1+1}^*(x_{h_2}) \dots p_{L_{t-1}+1}^*(x_{h_t}) \overline{G_{1L_1}} \cdot \overline{G_{L_1+1, L_2}} \cdot \dots \cdot \overline{G_{L_{t-1}+1, L_t}}$$

(here $\overline{G_{ab}} = G_{a, a+1} G_{a+1, a+2} \dots G_{b-1, b}$).

The last claim of the theorem is a consequence of the $\mathcal{S}_n$ -equivariant surjection

$$\mathbb{Q}[\mathcal{S}_n] \rightarrow E(L_*, H_*), \quad \sigma \mapsto \sigma \cdot \mu_{(L_*, H_*)}.$$

To prove a) it is enough to consider the action of the transpositions $\tau_i = (i, i+1)$ on the tree T containing the vertices $i, i+1$ or on the disjoint union of two trees, T'

and T'' , containing i and $i + 1$ respectively. In the case of one tree, the transform $\tau_i T$ is again a monotonic tree (and $\tau_i T \sim T$) if the path from 1 to $i + 1$ does not contain i . Otherwise the monotonic path $1 - \dots - h - i - (i + 1)$ is transformed into $1 - \dots - h - (i + 1) - i$ and the corresponding factor $G_{h,i+1}G_{i,i+1}$ should be replaced by $G_{h,i}G_{i,i+1} - G_{h,i}G_{h,i+1}$. The resulting monomials have monotonic trees of the same type with T . The case of two trees, i in T' and $i + 1$ in T'' , is simpler: $\tau_i(T' \sqcup T'')$ is a union of two monotonic trees ($h < i < j$ is equivalent to $h < i + 1 < j$) and obviously this union is of the same type with $T' \sqcup T''$.

It is enough to prove b) for a monotonic tree: we will show by induction on n that the bamboo $B_n = \mu_{(L_*= (n), H_*= (x_h))}$ generate the module $\text{Res}_{\mathcal{S}_{n-1}}^{\mathcal{S}_n} E_*^*(X, n)$. If σ is a permutation in $\mathcal{S}_{n-1}$, we denote by $\tilde{\sigma}$ its extension to $\mathcal{S}_n$: $\tilde{\sigma}(n) = n$. In the case $n = 3$ there is a unique monotonic tree which is not the monotonic bamboo B_3 , but this belongs to the $\mathcal{S}_2$ -orbit of B_3 :

$$\begin{array}{c} 2 \quad 1 \quad 3 \\ \hline \end{array} = \tau_1 \left(\begin{array}{c} 1 \quad 2 \quad 3 \\ \hline \end{array} \right) = \tau_1(B_3) \text{ and } \tau_1 \in \mathcal{S}_2.$$

Let us consider a monotonic tree T_n with n vertices and its monotonic subtree $T_{n-1} = T_n \setminus \{n\}$ (by monotonicity, the vertex n is connected to a unique other vertex, h). From the set of permutations $\pi \in \mathcal{S}_{n-1}$ with the property that $\pi(T_{n-1})$ is still monotonic we choose one such that $\pi(h) = j$ is maximal. Now we start a second induction on $n - j$, the number of vertices of $\pi(T_n)$ lying on the branches starting from j . If this number is equal to 1, then $j = n - 1$ and, by induction on n there are permutations σ_a in $\mathcal{S}_{n-2}$ and constants $c_a \in \mathbb{Q}$ such that

$$\pi(T_{n-1}) = \sum_a c_a \sigma_a(B_{n-1}).$$

The extension of a permutation $\tilde{\sigma}$ of a permutation in $\mathcal{S}_{n-2}$ does not change the edge $(n - 1) - n$ and we find $\tilde{\pi}(T_n) = \sum_a c_a \tilde{\sigma}_a(B_n)$ and therefore

$$T_n = \sum_a c_a \widetilde{\pi^{-1} \sigma_a}(B_n), \text{ where } \pi^{-1} \sigma_a \in \mathcal{S}_{n-1}.$$

If j is less than $n - 1$, then $j + 1$ is connected to j (by the maximality condition); applying the transposition τ_j , we obtain a non-monotonic tree $\tau_j \pi(T_n)$ (if $j \neq 1$). The sequence $k - (j + 1) - j$ should be replaced: we obtain a difference of monotonic trees $T'_n - T''_n$, in each of them n is connected with $j + 1$. The second induction will give two expansions

$$T'_n = \sum_p c'_p \sigma'_p(B_n), \quad T''_n = \sum_q c''_q \sigma''_q(B_n)$$

with $\sigma'_p, \sigma''_q \in \mathcal{S}_{n-1}$. Therefore T_n is in $\mathbb{Q}[\mathcal{S}_{n-1}](B_n)$:

$$T_n = \sum_p c'_p \widetilde{\pi^{-1} \tau_j \sigma'_p}(B_n) - \sum_q c''_q \widetilde{\pi^{-1} \tau_j \sigma''_q}(B_n).$$

In the case $j = 1$ the transposition τ_1 transforms $\pi(T_n)$ into a monotonic tree where n is connected with 2 and we can apply the second induction. $\square$

Corollary 2.14. *The $\mathcal{S}_{n-1}$ orbit of the monomial $x_h \otimes 1 \otimes \dots \otimes 1 G_{12} G_{23} \dots G_{n-1,n}$ coincides with $E_{n-1}^{(n-1)(2m-1)+|x_h|}(n, h)$ and*

$$\text{Res}_{\mathcal{S}_{n-1}}^{\mathcal{S}_n} E_{n-1}^{(n-1)(2m-1)+|x_h|}(n, h) \cong \mathbb{Q}[\mathcal{S}_{n-1}].$$

Proof. In the proof of the theorem we obtained a surjective $\mathcal{S}_{n-1}$ -map

$$\mathbb{Q}[\mathcal{S}_{n-1}] \rightarrow \text{Res}_{\mathcal{S}_{n-1}}^{\mathcal{S}_n} E_{n-1}^{(n-1)(2m-1)+|x_h|}(n, h), \quad \sigma \mapsto \sigma \cdot B_n.$$

On the other hand the number of monotonic trees with n vertices is $(n-1)!$. $\square$

3. THE $\mathcal{S}_n$ -MODULE $E_q^k(L_*, H_*)$

In this section we will study the symmetric structure of the modules $E_q^k(L_*, H_*)$ using two methods. We extend the main result of Lehrer and Solomon [LeSo] and we will give a general formula, but in an implicit form; next, in some particular cases, we will present explicit computations of the irreducible components and their multiplicities, using direct methods.

The symmetric structure of the bottom horizontal line is completely elementary: the types are given by $L_* = (1^{(n)})$, $H_* = (h_1^{(m_1)}, h_2^{(m_2)}, \dots, h_t^{(m_t)})$

$$\begin{array}{ccccccc} x_{h_1} & & x_{h_1} & & & & x_{h_t} & & x_{h_t} \\ \bullet & & \bullet & \dots & \dots & \dots & \bullet & & \bullet \\ 1 & & 2 & & & & n-1 & & n \\ \underbrace{\hspace{1.5cm}}_{m_1} & & & & & & \underbrace{\hspace{1.5cm}}_{m_t} \end{array}$$

(here $m_1, m_2, \dots, m_t$ are multiplicities of the elements $x_{h_1} \succ x_{h_2} \succ \dots \succ x_{h_t}$ in $\mathcal{B}$ and $m_1 + \dots + m_t = n$).

Example 3.1. In the case of distinct marks (all the multiplicities m_i are equal to 1) we obtain the largest possible “type” submodule:

$$E_0^{|H_*|}(1^{(n)}, (h_1, h_2, \dots, h_n)) \cong \mathbb{Q}[\mathcal{S}_n].$$

Proof. As $\sigma^{-1}(x_1 \otimes \dots \otimes x_n) = \pm x_{\sigma(1)} \otimes \dots \otimes x_{\sigma(n)}$, the character of this module is given by $\chi_{E_0^{|H_*|}}(\sigma) = \begin{cases} n! & \sigma = id \\ 0 & \sigma \neq id. \end{cases}$ $\square$

Example 3.2. In the case of a unique mark ($m_1 = n$)

$$E_0^{n|x_h|}(1^{(n)}, h^{(n)}) \cong \begin{cases} V(n) & \text{if } |x_h| \text{ is even or } n = 1 \\ V(1, 1, \dots, 1) & \text{if } |x_h| \text{ is odd and } n \geq 2. \end{cases}$$

Proof. This is the consequence of the relation

$$\tau_i(x_h \otimes \dots \otimes x_h) = (-1)^{|x_h|}(x_h \otimes \dots \otimes x_h).$$

$\square$

For the general type corresponding to the discrete graph we will use the notation

$$V^{\epsilon(m, h)} = \begin{cases} V(m) & \text{if } m = 1 \text{ or } |x_h| \text{ is even} \\ V(1, 1, \dots, 1) & \text{if } m \geq 2 \text{ and } |x_h| \text{ is odd.} \end{cases}$$


Proposition 3.3. The $\mathcal{S}_n$ structure of the type (L_*, H_*) , where $L_* = (1^{(n)})$, $H_* = (h_1^{(m_1)}, h_2^{(m_2)}, \dots, h_t^{(m_t)})$ is given by

$$E_0^{|H_*|}(1^{(n)}, H_*) \cong \text{Ind}_{\mathcal{S}_{m_1} \times \mathcal{S}_{m_2} \times \dots \times \mathcal{S}_{m_t}}^{\mathcal{S}_n} V^{\epsilon(m_1, h_1)} \otimes V^{\epsilon(m_2, h_2)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}.$$

Proof. The symmetric group $\mathcal{S}_n$ acts transitively on the set of 1-dimensional spaces $\mathbb{Q}\langle x_1 \otimes \dots \otimes x_n \rangle$, where m_1 factors (on different positions) coincide with x_{h_1} , ..., m_t factors coincide with x_{h_t} . The subgroup leaving invariant the 1-dimensional

subspace $\underbrace{x_{h_1} \otimes \dots \otimes x_{h_1}}_{m_1} \otimes \dots \otimes \underbrace{x_{h_t} \otimes \dots \otimes x_{h_t}}_{m_t}$ is the direct product $\mathcal{S}_{m_1} \times \mathcal{S}_{m_2} \times \dots \times \mathcal{S}_{m_t}$ (with the obvious notation: $\mathcal{S}_{m_1}$ acts on the subset $\{1, 2, \dots, m_1\}$, $\mathcal{S}_{m_2}$ acts on the subset $\{m_1 + 1, m_1 + 2, \dots, m_1 + m_2\}$, and so on) and the corresponding representation of this subgroup is $V^{\epsilon(m_1, h_1)} \otimes V^{\epsilon(m_2, h_2)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}$. General facts from the theory of induced representations (see for instance [Se], chapter 7) imply the result. $\square$

On the next horizontal line the types are given by a unique graph:



(here $x_{h_2} \succ \dots \succ x_{h_t}$, but h_1 could be equal to one of $h_2, \dots, h_t$ and we have $2 + m_2 + \dots + m_t = n$).

Proposition 3.4. *The $\mathcal{S}_n$ structure corresponding to the type $L_* = (2, 1^{(n-2)})$, $H_* = (h_1; h_2^{(m_2)}, \dots, h_t^{(m_t)})$ is given by*


$$E_1^{2m-1+|H_*|}(L_*, H_*) \cong \text{Ind}_{\mathcal{S}_2 \times \mathcal{S}_{m_2} \times \dots \times \mathcal{S}_{m_t}}^{\mathcal{S}_n} V(2) \otimes V^{\epsilon(m_2, h_2)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}.$$

Proof. This is similar to the previous proof: the module $E_*^*(L_*, H_*)$ is the direct sum of 1-dimensional subspaces $\mathbb{Q}\langle x_1 \otimes \dots \otimes x_n G_{ij} \rangle$, where on the i -th position lies x_{h_1} , on the j -th position is 1, and x_{h_2} is lying on m_2 (arbitrary) positions, $\dots, x_{h_t}$ on m_t positions, and these subspaces are permuted by $\mathcal{S}_n$. The subgroup leaving invariant the line $\mathbb{Q}\langle x_{h_1} \otimes 1 \otimes \underbrace{x_{h_2} \otimes x_{h_2} \otimes \dots \otimes x_{h_2}}_{m_2} \otimes \dots \otimes \underbrace{x_{h_t} \otimes \dots \otimes x_{h_t}}_{m_t} G_{12} \rangle$ is the product


$\mathcal{S}_2 \times \mathcal{S}_{m_2} \times \dots \times \mathcal{S}_{m_t}$ and the representation on this line is equivalent with $V(2) \otimes V^{\epsilon(m_2, h_2)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}$, hence the result. $\square$

The following types appear on the third horizontal line:


a) $L_* = (3, 1^{(n-3)})$, $H_* = (h_1, h_2^{(m_2)}, \dots, h_t^{(m_t)})$



b) $L_* = (2^{(2)}, 1^{(n-4)})$, $H_* = (h_1^{(2)}, h_2^{(m_2)}, \dots, h_t^{(m_t)})$



c) $L_* = (2^{(2)}, 1^{(n-4)})$, $H_* = (h_1, h_2, h_3^{(m_3)}, \dots, h_t^{(m_t)})$



(In the first two cases $h_2, \dots, h_t$ are distinct and h_1 could be one of them, in the third case $x_{h_1} \succ x_{h_2}, x_{h_3} \succ \dots \succ x_{h_t}$ and h_1 or h_2 are not necessarily different from $h_3, \dots, h_t$).

Example 3.5. For $n \geq 6$, the module $E_2^{2(2m-1)}(L_* = (3, 1^{(n-3)}), H_* = (1, 1^{(n-3)}))$ has the stable decomposition

$$V(1)_n \oplus V(2)_n \oplus V(1, 1)_n \oplus V(2, 1)_n.$$

Proof. Computing directly the corresponding character, we find for an arbitrary permutation $\sigma \in \mathcal{S}_n$ of type $(i_1; i_2; \dots; i_n)$ (here i_q is the number of cycles of length q): for any triple of 1-cycles $(i)(j)(k)$, $1 \leq i < j < k \leq n$, σ fixes the monomials $G_{ij}G_{ik}$ and $G_{ij}G_{jk}$ and this gives $2\binom{i_1}{3}$ such monomials. The 3-cycle (i, j, k) changes the sign of $G_{ij}G_{jk}$ and there is no other combination of cycles leaving $\mathbb{Q}\langle G_{ij}G_{jk} \rangle$ or $\mathbb{Q}\langle G_{ij}G_{ik} \rangle$ invariant. Therefore the character is

$$\chi_{E_2^{2(2m-1)}(\bullet-\bullet-\bullet \dots \bullet)}(i_1; \dots; i_n) = 2\binom{i_1}{3} - i_3.$$

Using Frobenius formula we obtain the next results:

	$\chi_V(i_1; \dots; i_n)$
$V(1)_n$	$i_1 - 1$
$V(1, 1)_n$	$\binom{i_1-1}{2} - i_2$
$V(2)_n$	$\frac{1}{2}i_1(i_1 - 3) + i_2$
$V(3)_n$	$\frac{1}{6}i_1(i_1 - 1)(i_1 - 5) + i_2(i_1 - 1) + i_3$
$V(2, 1)_n$	$\frac{1}{3}i_1(i_1 - 2)(i_1 - 4) - i_3$
$V(3, 1)_n$	$\frac{1}{8}i_1(i_1 - 1)(i_1 - 3)(i_1 - 6) + i_2\binom{i_1-1}{2} - \binom{i_2}{2} - i_4$

and as a consequence the above decomposition. $\square$

In the unstable cases, $n = 3, 4, 5$, this module decomposes as

$$\begin{aligned} n = 3: & \quad V(2, 1), \\ n = 4: & \quad V(3, 1) \oplus V(2, 2) \oplus V(2, 1, 1), \\ n = 5: & \quad V(4, 1) \oplus V(3, 2) \oplus V(3, 1, 1). \end{aligned}$$

The same decompositions appear if H_* is replaced by $H_* = (h_1, h_2^{(n-3)})$ with $|x_{h_1}|, |x_{h_2}|$ even.

Example 3.6. The module $E_2^{2(2m-1)}(L_* = (2^{(2)}, 1^{(n-4)}), H_* = (1^{(2)}; 1^{(n-4)}))$, for $n \geq 7$, has the stable decomposition

$$V(1)_n \oplus V(2)_n \oplus V(1, 1)_n \oplus V(3)_n \oplus V(2, 1)_n \oplus V(3, 1)_n.$$

Proof. Direct computation of the character gives non zero contributions only for factors of the form $(i)(j)(k)(l)$, $(i)(j)(k, l)$, $(i, j)(k, l)$ and (i, j, k, l) ; the result is

$$\chi_{E_*^*(\bullet-\bullet-\bullet-\bullet \dots \bullet)}(i_1; \dots; i_n) = 3\binom{i_1}{4} + i_2\binom{i_1}{2} - \binom{i_2}{2} - i_4$$

and from the same table we obtain the above decomposition. $\square$

In the unstable cases we obtain the decompositions

$$\begin{aligned} n = 4: & \quad V(3, 1), \\ n = 5: & \quad V(4, 1) \oplus V(3, 2) \oplus V(3, 1, 1) \oplus V(2, 2, 1), \\ n = 6: & \quad V(5, 1) \oplus V(4, 2) \oplus V(4, 1, 1) \oplus V(3, 3) \oplus V(3, 2, 1). \end{aligned}$$

Example 3.7. In the case $n = 4$ we have the decompositions

$$E_2^{2(2m-1)+2|x_h|}(\bullet_1^{x_h} - \bullet_2^{x_h} - \bullet_3^{x_h} - \bullet_4^{x_h}) \cong \begin{cases} V(3, 1) & \text{if } |x_h| = \text{even} \\ V(4) \oplus V(2, 2) & \text{if } |x_h| = \text{odd}. \end{cases}$$

Proof. The even case is the same as in the previous example; for the odd case direct computation of the character gives

σ	id	(12)	(123)	(1234)	$(12)(34)$
$\chi(\sigma)$	3	1	0	1	3

For instance:

$$\begin{aligned} (1234)p_1^*(x_h)p_2^*(x_h)G_{13}G_{24} &= p_2^*(x_h)p_3^*(x_h)G_{24}G_{13} = \\ &= -p_2^*(x_h)p_3^*(x_h)G_{13}G_{24} = -p_2^*(x_h)p_1^*(x_h)G_{13}G_{24} = \\ &= (-1)^{|x_h|+1}p_1^*(x_h)p_2^*(x_h)G_{13}G_{24}. \end{aligned}$$

□

Using some of these particular cases and the same proof as in Proposition 3.3 and 3.4, we obtain:

Proposition 3.8. The $\mathcal{S}_n$ structure of the modules corresponding to the types on the third horizontal line is given by:

$$\begin{aligned} a) E_2^*(L_* = (3, 1^{(n-3)}), H_* = (h_1, h_2^{(m_2)}, \dots, h_t^{(m_t)})) &\cong \\ \cong Ind_{\mathcal{S}_3 \times \mathcal{S}_{m_2} \times \dots \times \mathcal{S}_{m_t}}(V(2, 1) \otimes V^{\epsilon(m_2, h_2)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}); \\ b) E_2^*(L_* = (2^{(2)}, 1^{(n-4)}), H_* = (h_1^{(2)}, h_2^{(m_2)}, \dots, h_t^{(m_t)})) &\cong \\ \begin{cases} Ind_{\mathcal{S}_4 \times \mathcal{S}_{m_2} \times \dots \times \mathcal{S}_{m_t}}(V(3, 1) \otimes V^{\epsilon(m_2, h_2)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}) & |x_{h_1}| = \text{even}; \\ Ind_{\mathcal{S}_4 \times \mathcal{S}_{m_2} \times \dots \times \mathcal{S}_{m_t}}((V(4) \oplus V(2, 2)) \otimes V^{\epsilon(m_2, h_2)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}) & |x_{h_1}| = \text{odd}; \end{cases} \\ c) E_2^*(L_* = (2^{(2)}, 1^{(n-4)}), H_* = (h_1, h_2, h_3^{(m_3)}, \dots, h_t^{(m_t)})) &\cong \\ \cong Ind_{\mathcal{S}_2 \times \mathcal{S}_2 \times \mathcal{S}_{m_3} \times \dots \times \mathcal{S}_{m_t}}(V(2) \otimes V(2) \otimes V^{\epsilon(m_3, h_3)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}) \\ \cong Ind_{\mathcal{S}_4 \times \mathcal{S}_{m_3} \times \dots \times \mathcal{S}_{m_t}}((V(4) \oplus V(3, 1) \oplus V(2, 2)) \otimes V^{\epsilon(m_3, h_3)} \otimes \dots \otimes V^{\epsilon(m_t, h_t)}); \end{aligned}$$

Now we will analyze the top horizontal line; we start with the upper-left vertex of the trapezoid, $E_{n-1}^{(n-1)(2m-1)}(X, n)$.

Example 3.9. For small values of n the structure of this module is a consequence of the absence (for $n \geq 2$) of the one dimensional submodules of $E_{n-1}^*(X, n)$ and of the result of Section 2: $Res_{\mathcal{S}_{n-1}}^{\mathcal{S}_n} E_{n-1}^{(n-1)(2m-1)}(X, n) \cong \mathbb{Q}[\mathcal{S}_{n-1}]$:

$$\begin{aligned} n = 2: & E_1^{2m-1} \cong V(2); \\ n = 3: & E_2^{2(2m-1)} \cong V(2, 1); \\ n = 4: & E_3^{3(2m-1)} \cong V(3, 1) \oplus V(2, 1, 1); \\ n = 5: & E_4^{4(2m-1)} \cong V(4, 1) \oplus V(3, 2) \oplus V(3, 1, 1) \oplus V(2, 2, 1) \oplus V(2, 1, 1, 1). \end{aligned}$$

By Poincaré duality, the right side of the trapezoid has the same structure.

Example 3.10. For $n = 6$ there are seven $\mathcal{S}_6$ -modules without one dimensional submodules and satisfying $Res_{\mathcal{S}_5}^{\mathcal{S}_6} E_5^{5(2m-1)} \cong \mathbb{Q}[\mathcal{S}_5]$. A direct combinatorial computation of the character of $E_5^{5(2m-1)}$ will give its non zero values


σ	id	(123456)	$(123)(456)$	$(12)(34)(56)$
$\chi(\sigma)$	120	-1	-3	8

and therefore the next (asymmetric) decomposition: $E_5^{5(2m-1)} \cong$

$$\begin{aligned} & V(5, 1) \oplus 2V(4, 2) \oplus V(4, 1, 1) \oplus \dots \oplus 3V(3, 2, 1) \\ & V(2, 1, 1, 1, 1) \oplus V(2, 2, 1, 1) \oplus 2V(3, 1, 1, 1) \oplus V(2, 2, 2). \end{aligned}$$

One can find a character table of $\mathcal{S}_6$ in [L].

Alternatively, we can use the next general result of Stanley [St] and Lehrer-Solomon [LeSo] and the Frobenius reciprocity formula. Let us remind some notation necessary to present Lehrer-Solomon theorem: consider the subgroup generated by the n -cycle $c_n = (1, 2, \dots, n)$, $\langle c_n \rangle$, the elementary character φ_n of this cyclic group, $\varphi_n(c_n) = e^{\frac{2\pi i}{n}}$ and ε_n , the sign character of $\mathcal{S}_n$. More generally, for a partition $L_* \vdash n$, $L_* = (\lambda_1 \geq \lambda_2 \geq \dots \lambda_t \geq 1)$, let us denote by c_{L_*} the product of cycles $c_{\lambda_1} c_{\lambda_2} \dots c_{\lambda_t}$, $c_{\lambda_1} = (1, 2, \dots, L_1)$, $c_{\lambda_2} = (L_1 + 1, L_1 + 2, \dots, L_2), \dots, c_{\lambda_t} = (L_{t-1} + 1, L_{t-1} + 2, \dots, L_t)$. Associated to L_* we have the next diagram of groups and characters of one-dimensional representations:



where $\langle c_{L_*} \rangle$ is the subgroup generated by c_{L_*} , $\mathcal{S}_{L_*}$ is the direct product

$$\mathcal{S}_{L_*} \cong \mathcal{S}_{\lambda_1} \times \mathcal{S}_{\lambda_2} \times \dots \times \mathcal{S}_{\lambda_t},$$

C_{L_*} is the centralizer of c_{L_*} in $\mathcal{S}_{L_*}$:

$$C_{L_*} = \langle c_{\lambda_1} \rangle \times \langle c_{\lambda_2} \rangle \times \dots \times \langle c_{\lambda_t} \rangle.$$

The group N_{L_*} is generated by the elements v_i , “block transpositions” corresponding to equal parts $\lambda_i = \lambda_{i+1}$ in the partition L_* :

$$v_i = (L_{i-1} + 1, L_i + 1)(L_{i-1} + 2, L_i + 2) \dots (L_i = L_{i-1} + \lambda_i, L_{i+1} = L_i + \lambda_{i+1}).$$

The last group, Z_{L_*} , is the centralizer of c_{L_*} in $\mathcal{S}_n$ and it is a semi direct product

$$Z_{L_*} = C_{L_*} \rtimes N_{L_*}.$$

Lehrer-Solomon main formula involves the next characters:

$$\begin{aligned} \varphi_{L_*} : C_{L_*} &\rightarrow \mathbb{C}^*, & \varphi_{L_*} &= (\varphi_{\lambda_1} \otimes \dots \otimes \varphi_{\lambda_t}) \cdot \varepsilon_n|_{C_{L_*}}, \\ \alpha_{L_*} : N_{L_*} &\rightarrow \mathbb{C}^*, & \alpha_{L_*}(v_i) &= (-1)^{\lambda_i+1}, \\ \xi_{L_*} : Z_{L_*} &\rightarrow \mathbb{C}^*, & \xi_{L_*} &= \alpha_{L_*} \cdot \varphi_{L_*}. \end{aligned}$$

Theorem 3.11. (Stanley [St], Lehrer-Solomon [LeSo]) *The representation of $\mathcal{S}_n$ on the top component $\mathcal{A}^{n-1}(n)$ of the Arnold algebra has the character:*

$$\chi_{\mathcal{A}^{n-1}(n)} = \varepsilon_n \text{Ind}_{\langle c_n \rangle}^{\mathcal{S}_n}(\varphi_n).$$

On the upper horizontal line the types are parameterized by the n -monotonic bamboo and the monomials from the fixed basis $\mathcal{B}$:

$$\begin{array}{c} x_h \\ \bullet \text{---} \bullet \text{---} \dots \text{---} \bullet \\ 1 \quad 2 \quad \dots \quad n \end{array}$$

Proposition 3.12. *The $\mathcal{S}_n$ structure of the top component is given by*

$$\begin{aligned} \chi_{E_{n-1}^{(n-1)(2m-1)+|x_h|}(n,h)} &\cong \varepsilon_n \text{Ind}_{\langle c_n \rangle}^{\mathcal{S}_n}(\varphi_n), \\ \chi_{E_{n-1}^{(n-1)(2m-1)+i}(X,n)} &\cong \beta_i \cdot \varepsilon_n \text{Ind}_{\langle c_n \rangle}^{\mathcal{S}_n}(\varphi_n) \end{aligned}$$

(β_i is the i -th Betti number).

Proof. As the action of the symmetric group does not change the coefficients

$$\sigma(x_h \otimes 1 \otimes \dots \otimes 1_{G_{I_* J_*}}) = p_{\sigma(1)}^*(x_h) \cdot \sigma(G_{I_* J_*}) = p_1^*(x_h) \cdot \sigma(G_{I_* J_*})$$

we have the $\mathcal{S}_n$ -decomposition:

$$E_{n-1}^{(n-1)(2m-1)+i}(X, n) \cong \bigoplus_{x_h \in \mathcal{B} \cap H^i(X)} p_1^*(x_h) \cdot \mathcal{A}^{n-1}(n).$$

□

Example 3.13. *If the number of points is an odd prime number p , then the multiplicities m_λ of the irreducible $\mathcal{S}_p$ -modules are given by*

$$E_{p-1}^{(p-1)(2m-1)+|x_h|}(p, h) \cong \bigoplus_{\lambda \vdash p} m_\lambda V(\lambda),$$

$$m_\lambda = \frac{1}{p}(\dim V(\lambda) - \chi_\lambda(c_p)).$$

Proof. From the Frobenius reciprocity formula we obtain the expansion (here $V(\lambda^\varepsilon) = V(1, 1, \dots, 1) \otimes V(\lambda)$ with character $\chi_{\lambda^\varepsilon}$)

$$\begin{aligned} m_\lambda &= \langle \chi_V, \varepsilon_p \text{Ind}_{\langle c_p \rangle}^{\mathcal{S}_p}(\varphi_p) \rangle_{\mathcal{S}_p} = \langle \chi_{\lambda^\varepsilon}, \text{Ind}_{\langle c_p \rangle}^{\mathcal{S}_p}(\varphi_p) \rangle_{\mathcal{S}_p} = \\ &= \langle \text{Res}_{\langle c_p \rangle}^{\mathcal{S}_p} \chi_{\lambda^\varepsilon}, (\varphi_p) \rangle_{\mathbb{Z}_p} = \frac{1}{p} \sum_{k=0}^{p-1} \chi_{\lambda^\varepsilon}(c_p^k) e^{\frac{2k\pi i}{p}} \end{aligned}$$

in which all the values $\chi_{\lambda^\varepsilon}(c_p^k)$ are equal but not the first one:

$$\chi_{\lambda^\varepsilon}(c_p^0) = \dim V(\lambda^\varepsilon) = \dim V(\lambda).$$

□

Now we consider the type $L_* = (\lambda_1, \lambda_2, \dots, \lambda_t)$, $H_* = (1^{(t)})$ ($|L_*| = \sum(\lambda_i - 1)$). Translated into the language of types, the main formula of Lehrer-Solomon is

Theorem 3.14. (Lehrer-Solomon [LeSo]) *The representation of $\mathcal{S}_n$ on the component $E_{|L_*|}^{|L_*|(2m-1)}(L_*, 1^{(t)})$ has the character*

$$\chi_{E_{|L_*|}^{|L_*|(2m-1)}(L_*, 1^{(t)})} = \text{Ind}_{Z_{L_*}}^{\mathcal{S}_n}(\xi_{L_*}).$$

To extend this to a general type $L_* = (\lambda_1, \dots, \lambda_t)$, $H_* = (h_1, h_2, \dots, h_t)$, we modify the notation of Lehrer-Solomon as follows: the group $N_{(L_*, H_*)}$ is generated by the elements v_i corresponding to the transposition of equal marked bamboos: $\lambda_i = \lambda_{i+1}$, $h_i = h_{i+1}$ (remember that for equal lengths $\lambda_c = \lambda_{c+1} = \dots = \lambda_d$, the corresponding marks are decreasing, not necessarily strictly: $x_{h_c} \succeq x_{h_{c+1}} \succeq \dots \succeq x_{h_d}$); of course, v_i are given by the same product of disjoint transpositions.

The subgroup $Z_{(L_*, H_*)}$ is defined by the same formula:

$$Z_{(L_*, H_*)} = C_{L_*} \rtimes N_{(L_*, H_*)},$$

but now is, in general, smaller than the centralizer of c_{L_*} in $\mathcal{S}_n$. The character $\varphi_{L_*} : C_{L_*} \rightarrow \mathbb{C}^*$ is given by the same formula: for a permutation $\sigma \in \mathcal{S}_{L_*} = \mathcal{S}_{l_1} \times \mathcal{S}_{l_2} \times \dots \times \mathcal{S}_{l_t}$, the sign of $\sigma(x_{h_1} \otimes 1 \otimes \dots \otimes x_{h_2} \otimes 1 \otimes \dots \otimes x_{h_t} \otimes 1 \otimes \dots)$ is given only by the permutation of the exterior factors G_{ij} , like in Lehrer-Solomon definition of φ_{L_*} . The coefficients $\dots \otimes x_{h_i} \otimes 1 \otimes \dots \otimes x_{h_i} \otimes 1 \otimes \dots$ do have a contribution to the sign after the action of a permutation $\rho \in N_{(L_*, H_*)}$ if the degree $|x_{h_i}|$ is odd; therefore the character $\alpha_{(L_*, H_*)}$ should be modified as follows:

$$\alpha_{(L_*, H_*)} = \begin{cases} (-1)^{\lambda_i+1} & \text{if } |x_{h_i}| \text{ is even,} \\ (-1)^{\lambda_i} & \text{if } |x_{h_i}| \text{ is odd,} \end{cases}$$

and accordingly the character ξ is modified:

$$\xi_{(L_*, H_*)} = \varphi_{L_*} \cdot \alpha_{(L_*, H_*)}.$$

Finally we obtain the character of the Kriř algebra $E_*(X, n)$:

Theorem 3.15. *a) The $\mathcal{S}_n$ representation of the submodule $E_*(L_*, H_*)$ has the character*

$$\chi_{E_*(L_*, H_*)} = \text{Ind}_{Z_{(L_*, H_*)}}^{\mathcal{S}_n}(\xi_{(L_*, H_*)}).$$

b) The $\mathcal{S}_n$ representations of the component $E_q^k(X, n)$ has the character

$$\chi_{E_q^k(X, n)} = \sum_{\substack{|L_*|=q \\ |H_*|+q(2m-1)=k}} \text{Ind}_{Z_{(L_*, H_*)}}^{\mathcal{S}_n}(\xi_{(L_*, H_*)}).$$

c) The $\mathcal{S}_n$ representations of the Kriř algebra $E_*(X, n)$ has the character

$$\chi_{E_*(X, n)} = \sum_{(L_*, H_*)} \text{Ind}_{Z_{(L_*, H_*)}}^{\mathcal{S}_n}(\xi_{(L_*, H_*)}).$$

Proof. For a given partition $\Lambda_* = (\Lambda_1, \Lambda_2, \dots, \Lambda_t)$ of the set $\{1, 2, \dots, n\}$, where $|\Lambda_i| = \lambda_i$, let us denote by $E_*(\Lambda_*, H_*)$ the span of monomials associated to the marked graphs having Λ_* the set of connected components and H_* the corresponding marks. It is obvious that the symmetric group $\mathcal{S}_n$ acts transitively on the components of the direct sum

$$E_*(L_*, H_*) = \bigoplus_{\Lambda_*} E_*(\Lambda_*, H_*).$$

Now we fix the term $E_*(\Lambda_*, H_*)$ with

$$\Lambda_* = (\{1, \dots, L_1\}, \{L_1 + 1, \dots, L_2\}, \dots, \{L_{t-1} + 1, \dots, L_t\})$$

and we follow the proof of Lehrer-Solomon [LeSo] Section 4: the subgroup of $\mathcal{S}_n$ leaving $E_*(\Lambda_*, H_*)$ invariant is $Z_{(L_*, H_*)} = C_{L_*} \rtimes N_{(L_*, H_*)}$ (from the set v_i of permutations of connected components of equal size we have to consider only the

permutations of components with the same mark). The action of this subgroup onto $E_*^*(\Lambda_*, H_*)$ is a composition of:

C) the action of C_{L_*} on each component separately, and this is given componentwise by 3.12; its character is φ_{L_*} .

N) the action of $N_{(L_*, H_*)}$ which permutes identical marked trees; the rules of changing the sign were already explained.

The second and the third formulae of the theorem are direct consequences of the first one. $\square$

4. PROOFS OF PROPOSITION 1.2 AND PROPOSITION 1.3

In this section we will give a proof of Propositions 1.2 and 1.3. From the last section in [AAB] we will use the $\mathcal{S}_n$ -decomposition of $E_1^{2m-1}(X, n) \cong \mathcal{A}^1(n)$ and also the bases for the irreducible $\mathcal{S}_n$ -submodules described in this paper.

Proposition 4.1. [AAB] *The structure of the $\mathcal{S}_n$ -module $E_1^{2m-1}(X, n)$ is given by*

$$\begin{aligned} E_1^{2m-1}(X, 2) &\cong V(2), \\ E_1^{2m-1}(X, 3) &\cong V(3) \oplus V(2, 1), \\ E_1^{2m-1}(X, n) &\cong V(n) \oplus V(n-1, 1) \oplus V(n-2, 2) \quad \text{for } n \geq 4. \end{aligned}$$

We choose one nonzero element from each $\mathcal{S}_n$ -submodule:

$$\begin{aligned} G^n &= \sum_{i < j} G_{ij} \quad \text{in } V(n), \\ G_{12}^n &= \sum_{k \geq 3} (G_{1k} - G_{2k}) \quad \text{in } V(n-1, 1), \\ G_{1234} &= G_{14} - G_{13} + G_{23} - G_{24} \quad \text{in } V(n-2, 2) \end{aligned}$$

and by direct computation we find non-zero differentials:

Lemma 4.2. *The images of the elements G^n and G_{12}^n under the composition*

$$E_1^{2m-1} \xrightarrow{d} E_0^{2m} \xrightarrow{pr} \bigoplus_{i=1}^n p_i(H^{2m})$$

are given by

$$\begin{aligned} G^n &\mapsto (n-1) \sum_{i=1}^n p_i^*(w) \\ G_{12}^n &\mapsto (n-2)p_1^*(w) - (n-2)p_2^*(w). \end{aligned}$$

Lemma 4.3. *Let x and y be cohomology classes of positive degree (x in $\mathcal{B}$ and y in the dual bases $\mathcal{B}^*$) such that $xy = w$. The image of the element G_{1234} under the composition*

$$E_1^{2m-1} \xrightarrow{d} E_0^{2m} \xrightarrow{pr} \bigoplus_{i < j} p_i^*(\mathbb{Q}(x))p_j^*(\mathbb{Q}(y))$$

is given by

$$G_{1234} \mapsto p_1^*(x)p_4^*(y) - p_1^*(x)p_3^*(y) + p_2^*(x)p_3^*(y) - p_2^*(x)p_4^*(y).$$

Proposition 4.4. *If X is a projective manifold ($X \neq \mathbb{C}P^1$), then the “first” differential is injective:*

$$d : E_1^{2m-1}(X, n) \hookrightarrow E_0^{2m}(X, n).$$

Proof. Using twice the Schur lemma for the $\mathcal{S}_n$ -morphisms

$$V(n), V(n-1, 1) \hookrightarrow E_1^{2m-1}(X, n) \xrightarrow{d} E_0^{2m}(X, n)$$

these two submodules have isomorphic images through d in E_0^{2m} and trivial kernels because these morphisms are non-zero: $d(G^n) \neq 0$, $d(G_{12}^n) \neq 0$. If X is of complex dimension m greater than two, we can take in Lemma 4.3 x to be the Kähler class and $y = x^{m-1}$; if X is a projective curve, but not the projective line, the equation $xy = w$ has also non trivial solutions. $\square$

Remark 4.5. *In the remaining case, the differential*

$$d : E_1^1(\mathbb{CP}^1, n) \rightarrow E_0^2(\mathbb{CP}^1, n)$$

is injective only for $n = 2, 3$; for $n \geq 4$ we obtain

$$H_1^1(F(\mathbb{CP}^1, n)) \cong V(2)_n.$$

We will see in the last section that in the case of $\mathbb{CP}^1$, $n \geq 4$, the differential $E_q^q(\mathbb{CP}^1) \xrightarrow{d} E_{q-1}^{q+1}(\mathbb{CP}^1)$ has a nontrivial kernel for $q = 1, 2, \dots, n-3$ and it is injective for $q = n-2, n-1$.

Remark 4.6. *As a consequence of [LaSt], the Proposition 4.4 and the Proposition 1.2 are true for any formal space X whose rational cohomology satisfies Poincaré duality but not for cohomology spheres (for $n \geq 4$).*

Now we can give a proof of Proposition 1.2 by induction on q .

Proof. (of the Proposition 1.2) Let us suppose that the differential

$$d : E_{q-1}^{(q-1)(2m-1)} \rightarrow E_{q-2}^{(q-1)(2m-1)+1}$$

is injective and let $u \in E_q^{q(2m-1)}$ be a non zero cocycle. Let G_{ij} be the smallest exterior generator (in the reverse lexicographic order $G_{12} < G_{13} < G_{23} < G_{14} < \dots < G_{n-1, n}$) which appears in a non zero monomial (of the canonical basis) in u : $u = G_{ij}y + z$, $y \in E_{q-1}(G_{\alpha\beta} > G_{ij}) \setminus \{0\}$, $z \in E_q(G_{\alpha\beta} > G_{ij})$. In the right hand side of the equation

$$0 = d(u) = -G_{ij}dy + p_{ij}^*(\Delta)y + dz$$

the last two terms, $p_{ij}^*(\Delta)y$ and dz , the monomials in the canonical basis contain only factors $G_{\alpha\beta} > G_{ij}$; therefore $dy = 0$ by induction on q , and $y = 0$, and this gives a contradiction. $\square$

Proof. (of the Proposition 1.3) The monomials from the canonical basis lying on the top horizontal line are $p_1^*(x_h)G_{12}G_{i_2 3} \dots G_{i_{n-1} n}$, where $|x_h| = i \in [0, 2m]$ and $i_a \leq a$. The next composition is an isomorphism:

$$E_{n-1}^{(n-1)(2m-1)+i} \xrightarrow{d} E_{n-2}^{(n-1)(2m-1)+i+1} \xrightarrow{pr} \bigoplus_{\substack{I_*= (i_2, \dots, i_{n-1}) \\ i_a \leq a}} p_1^*(H^i)p_2^*(H^{2m})G_{i_2 3} \dots G_{i_{n-1} n}$$

$$p_1^*(x_h)G_{12}G_{i_2 3} \dots G_{i_{n-1} n} \mapsto p_1^*(x_h)p_2^*(w)G_{i_2 3} \dots G_{i_{n-1} n},$$

therefore the differential is injective on the top horizontal line. $\square$

5. AN ACYCLIC SUBALGEBRA OF THE KRIŘ MODEL

In [BMP] is introduced a quotient of the Kriř model, denoted by J_n , which is quasi isomorphic to $E_*(X, n)$, but the corresponding kernel is not $\mathcal{S}_n$ -stable. We will identify an acyclic subcomplex of $E_*(X, n)$, denoted by $E_*(w(X, n))$ (or simply by $E_*(w)$), which is also an $\mathcal{S}_n$ -submodule and a subalgebra, giving another smaller complex quasi isomorphic to the Kriř model:

$$SE_*(X, n) = E_*/E_*(w), \quad H^*(E_*(X, n)) \cong H^*(SE_*(X, n)).$$

The last isomorphism is now $\mathcal{S}_n$ -equivariant.

We start with a well known result in the theory of hyperplane arrangements, see for example [OT].

Definition 5.1. *The Arnold differential algebra $(\mathcal{A}^*(n), \partial)$ is defined by*

$$\mathcal{A}^*(n) = \bigwedge (G_{ij}, 1 \leq i < j \leq n) / (G_{ij}G_{ik} - G_{ij}G_{jk} + G_{ik}G_{jk})$$

(the generators G_{ij} have degree 1) and the differential of degree -1 is given by $\partial G_{ij} = 1$.

Proposition 5.2. *The Arnold algebra $(\mathcal{A}^*(n), \partial)$ is acyclic.*

Proof. Define the homotopy $h : \mathcal{A}^* \rightarrow \mathcal{A}^{*+1}$ by $h(\gamma) = G_{12}\gamma$ and verify that $\partial h + h\partial = id_{\mathcal{A}^*}$. $\square$

We denote by $E_*^{Top}(X, n)$ the submodule of the Kriř model $(E_*(X, n), d)$ given by the sum of the submodules of maximal total degree in each q -exterior degree

$$E_*^{Top}(X, n) = \bigoplus_{q=0}^{n-1} E_q^{Top}(X, n) = \bigoplus_{q=0}^{n-1} E_q^{2mn-q}(X, n).$$

It is obvious that $E_*^{Top}(X, n)$, the right side of the trapezoid, is a subcomplex and an ideal of $E_*(X, n)$.

Proposition 5.3. *There is an isomorphism of chain complexes of $\mathcal{S}_n$ -modules*

$$(\mathcal{A}^*(n), \partial) \cong (E_*^{Top}(X, n), d).$$

In particular

$$H^*(E_*^{Top}(X, n), d) = 0.$$

Proof. Using the standard basis $\{G_{I_*J_*} = G_{i_1j_1}G_{i_2j_2}\dots G_{i_qj_q}\}$ in $\mathcal{A}^q(n)$ and the basis $\{\prod_{h \notin J_*} p_h^*(w)G_{I_*J_*}\}$ in $E_q^{Top}(X, n)$ (here $2 \leq j_1 < j_2 < \dots < j_q \leq n, 1 \leq i_a < j_a$) we define the isomorphism

$$f : \mathcal{A}^q \rightarrow E_q^{Top} \text{ by } f(G_{I_*J_*}) = \prod_{h \notin J_*} p_h^*(w)G_{I_*J_*}.$$

Obviously f is $\mathcal{S}_n$ -equivariant (the degree of w is even) and f preserves the differentials:

$$\begin{aligned}
df(G_{I_*J_*}) &= d\left(\prod_{h \notin J_*} p_h^*(w) G_{I_*J_*}\right) \\
&= \sum_{a=1}^q (-1)^{a+1} \prod_{h \notin J_*} p_h^*(w) \cdot p_{i_a j_a}^*(\Delta) G_{i_1 j_1} \dots \widehat{G_{i_a j_a}} \dots G_{i_q j_q} \\
&= \sum_{a=1}^q (-1)^{a+1} \prod_{h \notin J_*} p_h^*(w) \cdot p_{j_a}^*(w) G_{i_1 j_1} \dots \widehat{G_{i_a j_a}} \dots G_{i_q j_q} \\
&= \sum_{a=1}^q (-1)^{a+1} \prod_{h \notin J_* \setminus \{j_a\}} p_h^*(w) G_{i_1 j_1} \dots \widehat{G_{i_a j_a}} \dots G_{i_q j_q} \\
&= f\left(\sum_{a=1}^q (-1)^{a+1} G_{i_1 j_1} \dots \widehat{G_{i_a j_a}} \dots G_{i_q j_q}\right) \\
&= f\partial(G_{I_*J_*}).
\end{aligned}$$

For the third equality we used $i_a \notin J_*$ and the equality

$$p_{i_a}^*(w) p_{i_a j_a}^*(\Delta) = p_{i_a}^*(w) p_{j_a}^*(w).$$

□

Proof. (of the Proposition 1.4) Now this is obvious. □

Now we will define three acyclic subcomplexes which generalize the previous subcomplex $(E_*^{Top}(X, n), d)$. For a fixed non empty subset $A \subset \{1, 2, \dots, n\}$ of cardinality $|A| = a \geq 2$ and a fixed sequence β of length $b = n - a$, $\beta = (x_1, x_2, \dots, x_b)$, where all the elements x_j belong to the fixed basis $\mathcal{B}$ and are different from w , we denote the increasing sequence of elements in $\{1, 2, \dots, n\} \setminus A$ by $b_1 < b_2 < \dots < b_b$, the product $\prod_{j=1}^b p_{b_j}^*(x_j)$ by $p^*(\beta)$, and its degree $\sum_{j=1}^b \deg(x_j)$ by $|\beta|$. Now we define subspace

$$E_*^{Top}(A, \beta) = \sum_{q=0}^{a-1} E_q^{2ma-q+|\beta|}(A, \beta)$$

by

$$E_q^{2ma-q+|\beta|}(A, \beta) = \mathbb{Q}\langle \prod_{i \in A \setminus J_*} p_i^*(w) p^*(\beta) G_{I_*J_*} \mid I_* \cup J_* \subset A, |J_*| = q \rangle$$

(in words: the scalars in the “complementary positions”, given by β , should be different from w , on the “forbidden positions”, corresponding to J_* , there is only 1, and all the other “possible positions” should be filled with the top class w).

Proposition 5.4. *For any A and β as before, the space $E_*^{Top}(A, \beta)$ is an acyclic subcomplex of the Križ model.*

Proof. By definition $E_*^{Top}(A, \beta)$ is the direct sum of its subspaces $E_q^{2ma-q+|\beta|}(A, \beta)$ and it is stable under the differential:

$$\begin{aligned}
d\left(\prod_{i \in A \setminus J_*} p_i^*(w) p^*(\beta) G_{I_*J_*}\right) &= \sum_{j_h \in J_*} \pm \prod_{i \in A \setminus J_*} p_i^*(w) p_{j_h}^*(w) p^*(\beta) G_{I_* \setminus \{i_h\} J_* \setminus \{j_h\}} = \\
&= \sum_{j_h \in J_*} \pm \prod_{i \in A \setminus (J_* \setminus \{j_h\})} p_i^*(w) p^*(\beta) G_{I_* \setminus \{i_h\} J_* \setminus \{j_h\}}.
\end{aligned}$$

This subcomplex is acyclic because of the isomorphism

$$(E_*^{Top}(A, \beta), d) \cong (\mathcal{A}^*(a), \partial)$$

given by

$$\prod_{i \in A \setminus J_*} p_i^*(w) p^*(\beta) G_{I_* J_*} \leftrightarrow (-1)^\beta G_{I_* J_*},$$

the differential of β is zero and the compatibility of the differentials d, ∂ was checked in the proof of 5.4. $\square$

Example 5.5. If $A = \{1, 2, \dots, n\}$, then β is the empty sequence, $p^*(\beta) = 1$ and $E_*^{Top}(\{1, 2, \dots, n\}, \phi) = E_*^{Top}(X, n)$.

Now we fix a number a from 2 to n and a sequence $\beta = (x_1, \dots, x_b)$ of length $b = n - a$ as before. We say that the sequence $\beta' = (y_1, \dots, y_b)$ is similar to β , $\beta \sim \beta'$, if there is a permutation $\sigma \in \mathcal{S}_b$ such that $y_i = x_{\sigma(i)}$ for $i = 1, 2, \dots, b$. We define a new subspace:

$$E_*^{Top}(a, \beta) = \sum_{\substack{|A|=a, \\ \beta' \sim \beta}} E_*^{Top}(A, \beta').$$

Proposition 5.6. For any number a and a sequence β as before, the space $E_*^{Top}(a, \beta)$ is an acyclic subcomplex and $\mathcal{S}_n$ -invariant.

Proof. The space is $\mathcal{S}_n$ -invariant by construction:

$$\sigma(E_*^{Top}(A, \beta)) = E_*^{Top}(\sigma A, \sigma \beta) = E_*^{Top}(\sigma A, \beta')$$

and the acyclicity is a consequence of the direct sum decomposition:

$$(E_*^{Top}(a, \beta), d) = \bigoplus_{\substack{|A|=a \\ \beta' \sim \beta}} (E_*^{Top}(A, \beta'), d).$$

$\square$

Finally, we take the whole collection of these subcomplexes:

$$E_*^*(w(X, n)) = \sum_{a=2}^n \sum_{\substack{\beta \text{ of length} \\ n-a}} E_*^{Top}(a, \beta).$$

Proposition 5.7. The space $E_*^*(w(X, n))$ is an acyclic, $\mathcal{S}_n$ -invariant subcomplex.

Proof. It is enough to show that the double sum is a direct sum: a monomial $x_1 \otimes \dots \otimes x_n G_{I_* J_*}$ from the canonical basis in $E_*^*(w(X, n))$ defines in a unique way the subset A and the factor β :

$$\begin{aligned} A &= \{i \in \{1, 2, \dots, n\} \mid x_i = w\} \cup J_*, \\ \beta &= (x_{h_1}, x_{h_2}, \dots, x_{h_b}), \end{aligned}$$

where $h_1 < h_2 < \dots < h_b$ are the elements of $\{1, 2, \dots, n\} \setminus A$. $\square$

Proposition 5.8. The projection map

$$E_*^*(X, n) \longrightarrow SE_*^*(X, n) = E_*^*(X, n) / E_*^*(w(X, n))$$

is a quasi-isomorphism.

Proof. This is obvious from the long exact sequence associated to

$$0 \rightarrow E_*^*(w(X, n)) \longrightarrow E_*^*(X, n) \longrightarrow SE_*^*(X, n) \rightarrow 0.$$

$\square$

6. AN EXAMPLE: $F(\mathbb{CP}^1, n)$

We analyze the cohomology algebra of the configuration space of the complex projective line using the symmetric structure of the Križ model. We encode the symmetric structure of a bigraded $\mathcal{S}_n$ -module H_*^* into the $\mathcal{S}_n$ -Poincaré polynomial:


$$SP_{H_*^*}(t, s) = \sum_{\lambda \vdash n} \left(\sum_{k, q} m_{q, \lambda}^k t^k s^q \right) V(\lambda),$$

where $m_{q, \lambda}^k$ is the multiplicity of the irreducible representation $V(\lambda)$ in the component H_q^k ; the double Poincaré polynomial of H_*^* is a consequence of $SP_{H_*^*}$: $P_{H_*^*}(t, s) = \sum_{k, q} \left(\sum_{\lambda \vdash n} m_{q, \lambda}^k \dim V(\lambda) \right) t^k s^q$.

For $n = 2$ and $n = 3$ we have the next tables of the symmetric group structure of the Križ model; using the injectivity properties of the differential, we obtain the first table and for the second table we have to use the vanishing of the cohomology on the left, top and the right side and also the acyclicity of the “interior part”

$$\bigoplus_{|A|=2} E_*^{Top}(A, 1) :$$

$$\begin{aligned} V(3) \oplus V(2, 1) &\cong \langle w \otimes 1 \otimes 1G_{12}, w \otimes 1 \otimes 1G_{13}, 1 \otimes w \otimes 1G_{23} \rangle \xrightarrow{d} \\ &\xrightarrow[\cong]{d} \langle w \otimes w \otimes 1, w \otimes 1 \otimes w, 1 \otimes w \otimes w \rangle : \end{aligned}$$



As a consequence we obtain

Lemma 6.1. *The non zero components of the cohomology algebra of $F(\mathbb{CP}^1, 2)$ and $F(\mathbb{CP}^1, 3)$ are*

$$\begin{aligned} H_0^0(F(\mathbb{CP}^1, 2)) &\cong V(2), & H_0^2(F(\mathbb{CP}^1, 2)) &\cong V(1, 1), \\ H_0^0(F(\mathbb{CP}^1, 3)) &\cong V(3), & H_1^3(F(\mathbb{CP}^1, 3)) &\cong V(3). \end{aligned}$$

In particular their symmetric Poincaré polynomials are

$$\begin{aligned} P_{F(\mathbb{CP}^1, 2)}(s, t) &= V(2) + t^2 V(1, 1), \\ P_{F(\mathbb{CP}^1, 3)}(s, t) &= (1 + st^3) V(3). \end{aligned}$$

Corollary 6.2. *The Poincaré polynomials of the unordered configuration spaces of the projective line are*

$$\begin{aligned} P_{C(\mathbb{CP}^1, 2)}(t) &= 1 \\ P_{C(\mathbb{CP}^1, 3)}(t) &= 1 + t^3. \end{aligned}$$

Another consequence of the last computation is the fact that the Serre spectral sequences of the fibrations

$$\mathcal{F}_n : F(\mathbb{C}, n-1) \hookrightarrow F(\mathbb{CP}^1, n) \rightarrow \mathbb{CP}^1$$

do not degenerate at $\mathbf{E}_2^{*,*}$ (for $n \geq 3$): using the vanishing of the first and second cohomology of $F(\mathbb{CP}^1, 3)$ and the projection $p : F(\mathbb{CP}^1, n) \rightarrow F(\mathbb{CP}^1, 3)$, we obtain the diagram (we use bold $\mathbf{E}_*^{*,*}$ for the components in the spectral sequences):

$$\begin{array}{ccccc} \mathbf{E}_2^{0,1}(\mathcal{F}_3) = \mathbb{Q}\langle G_{12} \rangle & \xrightarrow{p^*} & \mathbf{E}_2^{0,1}(\mathcal{F}_n) \cong \mathcal{A}^1(n-1) & & \\ & \searrow d_2 & & \searrow d_2 & \\ & \cong & \mathbf{E}_2^{2,0}(\mathcal{F}_3) \cong \mathbb{Q}\langle w \rangle & \xrightarrow[p^*]{\cong} & \mathbf{E}_2^{2,0}(\mathcal{F}_n) \cong \mathbb{Q}\langle w \rangle \end{array}$$

and we find that the differential d_2 is surjective for $n \geq 3$. These spectral sequences degenerate at E_3 : the two non zero columns are given by

$$\begin{aligned} \mathbf{E}_\infty^{0,*} &= \mathbf{E}_3^{0,*} = \ker d_2 \cong \mathcal{A}^*(n-1)/(G_{12}), \\ \mathbf{E}_\infty^{2,*} &= \mathbf{E}_3^{2,*} \cong \mathbf{E}_2^{2,*}/\text{Im } d_2 \cong \mathbb{Q}\langle wG_{12} \rangle \otimes \mathcal{A}^*(n-1)/(G_{12}). \end{aligned}$$

Proposition 6.3. [FZ] *The cohomology algebra of the configuration space $F(\mathbb{CP}^1, n)$ ($n \geq 3$) is given by*

$$H^*(F(\mathbb{CP}^1, n)) \cong H^*(F(\mathbb{CP}^1, 3)) \otimes \mathcal{A}^*(n-1)/(G_{12}).$$

In particular, its Poincaré polynomial is

$$P_{F(\mathbb{CP}^1, n)}(t) = (1+t^3)(1+2t)(1+3t)\dots(1+(n-2)t).$$

Using the results from section 5, we detect the nonzero bigraded components of the cohomology algebra and (partially) its $\mathcal{S}_n$ -structure.

Proof. (of the Theorem 1.5) The first cohomology group is

$$H^1(F(\mathbb{CP}^1, n)) = H_1^1 \cong V(n-2, 2),$$

and the subalgebra generated by degree 1 elements is contained in $\bigoplus_{q=0}^{n-3} H_q^q$. The element

$$\gamma = 2(n-2) \sum_{i < j} p_i^*(w)G_{ij} - \sum_{i < j} \sum_{k \neq i, j} p_k^*(w)G_{ij} \in E_1^3(\mathbb{CP}^1, n)$$

is a cocycle in the $V(n)$ -isotypic component. It can not be a coboundary because $V(n)$ is missing from E_2^2 :

$$\begin{aligned} E_2^2(\mathbb{CP}^1, n) &\cong \mathcal{A}^2(n) \cong \\ &\cong 2V(1)_n \oplus 2V(2)_n \oplus 2V(1, 1)_n \oplus V(3)_n \oplus 2V(2, 1)_n \oplus V(3, 1)_n \end{aligned}$$

(this is correct in the stable case $n \geq 7$, see [CF] or [AAB]; the trivial module $V(n)$ does not appear in the unstable cases either).

As $\beta_3 = 1 + \sum_{2 \leq i < j < k \leq n-2} ijk$ and the component H_3^3 contains a submodule of dimension $\beta_3 - 1$, we obtain

$$H_1^3(F(\mathbb{CP}^1, n)) \cong V(n).$$

The ideal generated by γ is contained in $\bigoplus_{q=1}^{n-2} H_q^{q+2}$ and algebra structure shows that all the other bigraded components are zero. $\square$

The module H_2^2 is a quotient of $E_2^2 \cong \mathcal{A}^2(n)$ (its decomposition into irreducible modules was given in the last proof) and also a quotient of

$$\begin{aligned} H_1^1 \wedge H_1^1 &\cong \bigwedge^2 V(2)_n \\ &\cong V(1, 1)_n \oplus V(2, 1)_n \oplus V(1, 1, 1)_n \oplus V(3, 1)_n \end{aligned}$$

(see [AAB]); the intersection of these decompositions gives (for $n \geq 7$) the inclusion

$$H_2^2 < V(1, 1)_n \oplus V(2, 1)_n \oplus V(3, 1)_n$$

and computing their dimensions this inclusion becomes an equality:

$$\begin{aligned} \beta_2 &= \sum_{2 \leq i < j \leq n-2} ij = \frac{(n-4)(n-3)(3n^2 - n + 2)}{24} \\ &= \frac{(n-1)(n-2)}{2} + \frac{n(n-1)(n-4)}{3} + \frac{4(n-1)(n-3)(n-6)}{8} \\ &= \dim V(1, 1)_n + \dim V(2, 1)_n + \dim V(3, 1)_n. \end{aligned}$$

Similar computations give the unstable cases of the next proposition.

Proposition 6.4. *The decomposition of the second cohomology group becomes stable for $n \geq 7$ and it is given by*

$$H^2(F(\mathbb{CP}^1, n)) = H_2^2 \cong V(1, 1)_n \oplus V(2, 1)_n \oplus V(3, 1)_n.$$

In the unstable cases we have

$$\begin{aligned} H^2(F(\mathbb{CP}^1, n)) &= 0 && \text{for } n = 2, 3, 4; \\ H^2(F(\mathbb{CP}^1, 5)) &\cong V(3, 1, 1) \\ H^2(F(\mathbb{CP}^1, 6)) &\cong V(4, 1, 1) \oplus V(3, 2, 1). \end{aligned}$$

REFERENCES

- [A] V.I. Arnold, *Cohomology of dyed braids*, Mathematical Notes, 5(2) (1969), 138-140.
- [AAB] S. Ashraf, H. Azam, B. Berceanu, *Representation stability of power sets and square free polynomials*, arXiv: 1106.4926v1 [math.RT] (2011).
- [AsB] S. Ashraf, B. Berceanu, *Equivariant Lefschetz structure of the Križ model*, in preparation.
- [AzB] H. Azam, B. Berceanu, *Cohomology of configuration spaces of Riemann surfaces*, in preparation.
- [Bg] G. Bergman. *The diamond lemma for ring theory*, Advances In Mathematics 29(2) (1978), 178-218.
- [BMP] B. Berceanu, M. Markl, S. Papadima, *Multiplicative models for configuration spaces of algebraic varieties*, Topology 44 (2005), 415-440.
- [Bz] R. Bezrukavnikov, *Koszul DG-algebras arising from configuration spaces*, Geometric and Functional Analysis, Vol 4, No. 2 (1994), 119-135.
- [CF] T. Church, B. Farb, *Representation Theory and Homological Stability*, arXiv:1008.1368v1 [math.RT] (2010).
- [CT1] F. Cohen, L. Taylor, *Configuration spaces: Applications to Gelfand- Fuks cohomology*, Bull. Amer. Math. Soc. 84 (1978), 134-136.

- [CT2] F. Cohen, L. Taylor, *Computations of Gelfand-Fuks cohomology, the cohomology of function spaces, and the cohomology of configuration spaces*; in: Geometric Applications of Homotopy Theory I, Proceedings, Evanston 1977; Lecture Notes in Mathematics vol. 657, 106-143, Springer-Verlag 1978.
- [DGMS] P. Deligne, P. Griffiths, J. W. Morgan, D. Sullivan, *The real homotopy type of Kähler manifolds*, Invent. Math. 29 (1975) 245-254.
- [FH] W. Fulton, J. Harris, *A First Course in Representation Theory*, Graduate Texts in Mathematics 129, Springer-Verlag Berlin, 1991.
- [FM] W. Fulton, R. MacPherson, *A compactification of configuration spaces*, Annals of Mathematics, Vol. 139, No. 1 (1994), 183-225.
- [FZ] E. M. Feichtner, G. M. Ziegler, *The integral cohomology algebras of ordered configuration spaces of spheres*, Documenta Mathematica 5 (2000), 115-139.
- [K] I. Križ, *On the rational homotopy type of configuration spaces*, Annals of Mathematics, Vol. 139, No. 2 (1994), 227-237.
- [L] W. Lederman, *Introduction to Group Characters*, Cambridge University Press, 1987.
- [LaSt] P. Lambrechts, D. Stanley, *A remarkable DG module model for configuration spaces*, Algebraic and Geometric Topology 8 (2008), 1191-1222.
- [LeSo] G. I. Lehrer, L. Solomon, *On the action of the symmetric group on the cohomology of the complement of its reflecting hyperplanes*, Journal of Algebra 104, (1986), 410-424.
- [OT] P. Orlik and H. Terao, *Arrangements of Hyperplanes*, Grundlehren der mathematische Wissenschaften, vol. 300. Springer-Verlag, Berlin, 1992.
- [St] R. P. Stanley, *Some aspects of groups acting on finite posets*, J. Combinatorial Theory, Series A, 32 (1982), 132-161.
- [Se] J. P. Serre, *Linear Representations of Finite Groups*, Graduate Texts in Mathematics 42, Springer-Verlag, Berlin, 1977.

¹ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES, GC UNIVERSITY, LAHORE-PAKISTAN.
E-mail address: samia.ashraf@yahoo.com

²ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES, GC UNIVERSITY, LAHORE-PAKISTAN.
E-mail address: centipedes.united@gmail.com

³ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES, GC UNIVERSITY, LAHORE-PAKISTAN,
AND INSTITUTE OF MATHEMATICS SIMION STOILOW, BUCHAREST-ROMANIA (PERMANENT ADDRESS).
E-mail address: Barbu.Berceanu@imar.ro